

# A Unique Find of the Ammonite *Kamerunoceras* (Acanthoceratidae, Ammonoidea) from the Turonian (Upper Cretaceous) of the South–Western Crimea

E. Yu. Baraboshkin<sup>a</sup>, \* (ORCID: 0000-0003-4373-1543) and P. A. Fokin<sup>a</sup>, \*\* (ORCID: 0009-0006-4303-3330)

<sup>a</sup> Moscow State University, Moscow, 119991 Russia

\*e-mail: EJBaraboshkin@mail.ru

\*\*e-mail: fokinpav@rambler.ru

Received November 7, 2023; revised December 19, 2023; accepted March 5, 2024

**Abstract**—A Turonian ammonite, *Kamerunoceras* sp. ex gr. *turonienne* (d’Orb.) has been found for the first time in the southwestern Crimea, in the section of the Aksu–Dere ravine, northward of the village of Kudrino (Kacha River basin). This is the first discovery of representatives of this genus in Russia.

**Keywords:** ammonite, *Kamerunoceras* sp. ex gr. *turonienne*, upper Cretaceous, Turonian, biostratigraphy, southwestern Crimea

DOI: 10.3103/S0145875224700212

## INTRODUCTION

Finds of ammonites are relatively rare in the Turonian deposits of Crimea and the vast majority of them originate from deposits of the upper Turonian Substage (Naidin et al., 1981; Alekseev, 1989; *Atlas melovoi fauny...*, 1997). The only mention of lower Turonian ammonites from sections of the Kacha–Bodrak interfluvium (members VIII and/or IX) is given in (Naidin et al., 1981): “subfamily Collignoniceratinae Wright et Wright (unidentifiable cast) and *Prionocyclus*? aff. *neptuni* (Geinitz).” Unfortunately, these finds are not depicted and are absent in the collections, which makes it impossible to evaluate them.

In 2023, P.A. Fokin found a large fragment of the ammonite *Kamerunoceras* sp. ex gr. *turonienne* (d’Orb.) in the reference section of Turonian deposits of the southwestern Crimea, in the Aksu–Dere ravine, during student practice. The find was made in a dirt road, on the rise from the ravine to the watershed, in Member IX of unevenly layered limestones with flint nodules, about 35 m above the well-known level of bituminous shale and top of the Cenomanian Stage (Figs. 1, 2). This is the first record of *Kamerunoceras* sp. ex gr. *turonienne* (d’Orb.) not only in the southwestern Crimea but also throughout Russia. Therefore, despite its poor preservation, it is of significant value and expands our knowledge of the distribution of representatives of this genus.

## PALEONTOLOGICAL DESCRIPTION

The ammonite was described using standard terminology and measurements of shells given in (*Atlas melovoi fauny...*, 1997). The collection is deposited in the Museum of Earth Science, Moscow State University under no. 149.

Order Ammonoidea Zittel, 1884

Suborder Ammonitina Hyatt, 1889

Superfamily Acanthoceratoidea de Grossouvre, 1894

Family Acanthoceratidae de Grossouvre, 1894

Subfamily Euomphaloceratinae Cooper, 1978

Genus *Kamerunoceras* Reymont, 1954

*Kamerunoceras* sp. ex gr. *turonienne* (d’Orbigny, 1850)

Figs. 3a, 3b

**Description.** The specimen is represented by approximately a quarter of a large whorl belonging to the living chamber. It is strongly flattened parallel to the symmetry plane and partially replaced by white and light gray flint. Shell is semi-evolute; its diameter was about 130 mm. The height of the whorl measured between the ribs is 50 mm. It is difficult to determine the cross-sectional shape. It is probably closer to the rectangular or square shape (Fig. 3b). Ribs are slightly curved backwards, rough, and simple. They begin from the umbilical seam almost unnoticeably and are ornamented with a umbilical tubercle or an elongated, laterally flattened swelling. Ribs are significantly weakened in the middle part of the sides and an elongated swelling appears towards the ventral side and




Fig. 1. Location of the section: (a) in Crimea (arrow), (b) in the Kacha River basin, to the north of the village of Kudrino (asterisk).

ends at the ventral bend with a slightly flattened internal ventrolateral tubercle. Another tubercle (external ventrolateral high and pointed tubercle) is located next to the previous one, but on the ventral side (Fig. 3b). The ventral side is almost invisible, which makes it difficult to establish the presence of a keel; in any case, it is hardly pronounced. There are seven ribs on a quarter of the whorl.

**Comparison.** The discovered specimen is close to representatives of different Turonian ammonite genera from different families. It resembles large whorls of some individuals of *Collignonicerias* Breistroffer, 1947 and *Prionocyclus* Meek, 1876 of the family Collignoniceratidae Wright et Wright, 1951; however, the latter have a well-developed keel and often additional thinner lateral ribs on the outer whorls. Among *Prionocyclus*, the coarse ribbed form of *Prionocyclus germari* (Reuss) is closest to our specimen; however, this species is recorded in the upper Turonian (Kennedy et al., 2001).

Among the early Turonian representatives of the subfamily Euomphaloceratinae Cooper, 1978, two genera, *Pseudaspidoceras* Hyatt, 1903 and *Kamerunoceras* Reymont, 1954, are closest to our specimen. In *Pseudaspidoceras*, ribs are almost always smoothed or there are thinner secondary, sometimes loop-like ribs on large whorls. Among the most coarsely ribbed forms, which have no additional ribs, are *Pseudaspidoceras paganum* Reymont and *Pseudaspidoceras pseudonodosoides* (Chof.); however, both of them have low external ventrolateral tubercles or have none of them at all. The pattern of ventrolateral tubercles is

most similar to our specimen only in *Kamerunoceras*. Among the representatives of this genus, *Kamerunoceras turoniense* (d'Orb.) (a zonal form that begins the middle Turonian sequence) is most similar to our specimen. *Kamerunoceras jacobsoni* Reymont is also a closely related form; however, it has equal ventrolateral tubercles, which is not observed in our specimen. Based on the above, we believe that the most accurate indication for the discovered ammonite is *Kamerunoceras* sp. ex gr. *turoniense* (d'Orbigny, 1850).


A detailed description of the species *Kamerunoceras turoniense* (d'Orb.), an image of the lectotype, and the discussion of the problems of its selection are given in (Kennedy and Wright, 1979). It contains a list of synonyms; supplemented synonyms for this species can be found in (Ayoub-Hannaa and Fürsich, 2012; Kennedy and Gale, 2020).

**Occurrence.** The upper part of the lower Turonian–lower part of the middle Turonian of England, France, Spain, Portugal, Morocco, Algeria, Tunisia, Lebanon, Egypt, Israel, Nigeria, Cameroon, Madagascar, Mexico, Texas and New Mexico (United States), Brazil, Colombia, Venezuela, Tajikistan (?), and Crimea (Fig. 4).


**Material.** One specimen, MZ MGU no. 149/7.

## DISCUSSION


In accordance with the decisions of the Colloquium on the Turonian Stage, a recommendation on the three-folder division of the Turonian (Robaszynski, 1983; Bengtson, 1996), including its



**Fig. 2.** Structure of the lower part of the lower–middle Turonian section of the Aksu-Dere ravine and distribution of macrofauna, including *Kamerunoceras* sp. ex gr. *turoniense* (d’Orb.) (our data). The boundaries of the zones of inocerams and planktonic foraminifers are indicated according to (Aleksiev, 1989; Kopaevich and Walaszczyk, 1990; Kopaevich and Walaszczyk, 1993) and the boundaries of the nannoplankton zones according to (Shcherbinina and Gavrilov, 2016).



**Fig. 3.** *Kamerunoceras* sp. ex gr. *turoniense* (d'Orb.), specimen no. 149/7: (a) lateral view, (b) reconstructed section of the shell through the main ribs; lower Turonian, Member IX, Aksu-Dere ravine, Bakhchisaray district, Republic of Crimea. Collected by Fokin, 2023. The specimen is covered with ammonium chloride. Photograph by E.Yu. Baraboshkin.



**Fig. 4.** Scheme of geographical distribution of *Kamerunoceras turoniense* (d'Orb.) on a paleogeographic basis (after R. Blakey, with changes).

division in the General Scale of the Cretaceous System (Olfer'ev and Alekseev, 2002), was accepted. It was proposed to establish the boundary by the first appearance of the ammonite *Collignoniceras woollgari* (Mantell) in the Rock Canyon section near Pueblo, Colorado, which is considered as a limitotype (GSSP) of this boundary (Bengtson, 1996). This level is almost consistent with the appearance of *Kamerunoceras turoniense* (d'Orb.), which was chosen as an index species of the same name subzone of the *Collignoniceras woollgari* zone for southern Europe (Amedro et al., 2020) and in the General Scale of the Upper Cretaceous of Russia (Olfer'ev and Alekseev, 2002), although it occurs in the upper part of the lower Turonian (Kennedy et al., 2015).

The Turonian deposits of the southwestern Crimea were long divided into two substages (Naidin et al., 1981; Alekseev, 1989). Later, the possibility of three-

folder division of the Turonian in the Aksu-Dere section was discussed based on microfauna and inoceram (Kopaevich and Walaszczyk, 1990; Kopaevich and Walaszczyk, 1993). The base of the Middle Turonian was conventionally defined along the lower boundary of Member X and the *Inoceramus lamarcki* Zone, about 18 m above the black shale horizon. In 2016, the boundary between the lower Turonian and middle Turonian in the Aksu-Dere section was established about 5 m above the bituminous shale horizon based on nanoplankton (Shcherbinina and Gavrillov, 2016, Fig. 2). At the same time, a certain difficulty of establishing this boundary was noted (Shcherbinina and Gavrillov, 2016, p. 294). According to our measurement of the thickness of the section, the base of Member X (as understood by (Alekseev, 1989)) and, consequently, the base of the *Inoceramus lamarcki* zone, is about 50 m above the black shale horizon and 15 m

above the level of the find of the ammonite (Fig. 2). If our measurements are sufficiently accurate and the information about the occurrence of *Inoceramus lamarcki* (Park.) is correct, the find of *Kamerunoceras* sp. ex gr. *turonienne* (d'Orb.) refers to the upper part of the lower Turonian rather than to the base of the middle Turonian. Obviously, this issue requires additional study of the microfauna.

### CONCLUSIONS

The specimen of *Kamerunoceras* sp. ex gr. *turonienne* (d'Orb.) is the first record of a representative of the genus *Kamerunoceras* in sections of Russia; it expands our knowledge of the distribution of this biostratigraphic marker. Although the species *Kamerunoceras turonienne* (d'Orb.) is an index of the same-name subzone of the *Collignoniceratites* zone of the middle Turonian base, its distribution interval is wider and its stratigraphic position in the Aksu-Dere section indicates the upper part of the lower Turonian.

### ACKNOWLEDGMENTS

The authors are grateful to V.V. Arkadiev (St. Petersburg State University) for constructive comments and editorial correction of the article.

### FUNDING

This study was supported by the Russian Science Foundation (no. 22-17-00091), <https://rscf.ru/project/22-17-00091/>.

### CONFLICT OF INTEREST

The authors of this work declare that they have no conflicts of interest.

### REFERENCES

- Alekseev, A.S., Upper Cretaceous, in *Geologicheskoe stroenie Kachinskogo podnyatiya Gornogo Kryma. Stratigrafiya mezozoya* (Geological Structure of the Kachinsky Uplift of the Mountain Crimea. Mesozoic Stratigraphy), Mazarovich, O.A. and Mileev, V.S., Eds., Moscow: Mosk. Gos. Univ., 1989.
- Amedro, F., Robaszynski, F., Chatelier, H., et al., Des nouveaux tuffeaux du Turonien moyen et sur leurs faunes d'ammonites en Touraine Meridionale new data on Middle Turonian tuffeaux of Southern Touraine and on their ammonites faunas, *Bul. Inf. Geol. Bass. Paris*, 2020, vol. 57, no. 2, pp. 3–33.
- Atlas melovoi fauny Yugo-Zapadnogo Kryma* (Atlas of Cretaceous fauna of South-Western Crimea), Arkadiev, V.V. and Bogdanov, T.N., Eds., St. Petersburg: Pangeya, 1997.
- Ayoub-Hannaa, W. and Fursich, F.T., Cenomanian-Turonian ammonites from eastern Sinai, Egypt, and their biostratigraphic significance, *Beringeria*, 2012, vol. 42, pp. 57–92.

- Bengtson, P., The Turonian Stage and Substage boundaries, *Bul. Inst. Roy. Sci. Nat. Belgique, Sci. Terre*, 1996, vol. 66, pp. 69–79.
- Kennedy, W.J. and Wright, C.W., On *Kamerunoceras* Reyment, 1954 (Cretaceous Ammonoidea), *J. Paleontol.*, 1979, vol. 53, no. 5, pp. 1165–1178.
- Kennedy, W.J. and Gale, A.S., The ammonite *Kamerunoceras* Reyment, 1954 from the Lower Turonian (Upper Cretaceous) of Goulmima, south-eastern Morocco, *N. Jahrb. Geol. Palaont. Abh.*, 2020, vol. 298, no. 2, pp. 197–202.
- Kennedy, W.J., Cobban, W.A., and Landman, N.H., A revision of the Turonian members of the ammonite subfamily Collignoniceratinae from the United States Western Interior and Gulf Coast, *Bull. Am. Mus. Nat. Hist.*, 2001, vol. 267.
- Kennedy, W.J., Bilotte, M., and Melchior, P., Turonian ammonite faunas from the Southern Corbieres, Aude, France, *Acta Geol. Pol.*, 2015, vol. 65, no. 4, pp. 437–494.
- Kopaevich, L.F. and Walaszczyk, I., An integrated inoceramid-foraminiferal biostratigraphy of the Turonian and Coniacian strata in South-Western Crimea, Soviet Union, *Acta Geol. Pol.*, 1990, vol. 40, nos. 1–2, pp. 83–95.
- Kopaevich, L.F. and Walaszczyk, I., Subdivision of Turonian–Coniacian beds in the Aksudere section based on inocerams and foraminifers, *Vestn. Mosk. Univ., Ser. 4: Geol.*, 1993, no. 5, pp. 70–82.
- Naidin, D.P., Alekseev, A.S., and Kopaevich, L.F., Fauna from Turonian deposits of the Kacha-Bodrak interfluvium (Crimea) and Cenomanian–Turonian boundary, in *Evolutsiya organizmov i biostratigrafiya serediny melovogo perioda* (Organic Evolution and Biostratigraphy in the mid-Cretaceous Period), Naidin, D.P. and Krasilov, V.A., Eds., Vladivostok: Dalnevost. Nauchn. Tsentr Akad. Nauk SSSR, 1981.
- Olfer'ev, A.G. and Alekseev, A.S., The Global Upper Cretaceous Scale, *Stratigr. Geol. Correl.*, 2002, vol. 10, no. 3, pp. 270–285.
- Robaszynski, F., Conclusions au Colloque sur le Turonien. Echeilles biostratigraphiques integrees, *Mem. Mus. Natl. Hist. Nat. Paris, Ser. C. Sci. Terre*, 1983, vol. 49, pp. 209–230.
- Shcherbinina, E.A. and Gavrillov, Yu.O., The nannoplankton zonation of Cenomanian–Santonian deposits of southwestern Crimea, in *Melovaya Sistema Rossii i blizhnego zarubezh'ya: problemy stratigrafii i paleogeografii: Sb. nauch. trudov* (Coll. Sci. Works “Cretaceous System of Russia and CIS Countries: Problems of Stratigraphy and Paleogeography”), Simferopol: Chernomorpres, pp. 292–294.
- Du Vivier, A.D.C., Jacobson, A.D., Lehn, G.O., et al., Ca isotope stratigraphy across the Cenomanian–Turonian OAE 2: links between volcanism, seawater geochemistry, and the carbonate fractionation factor, *Earth Planet. Sci. Lett.*, 2015, vol. 416, pp. 121–131.

Translated by D. Zabolotny

**Publisher's Note.** Allerton Press remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.